



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



*Published to advance the Science of cold-blooded vertebrates*

## SCOTCH SEA TROUT IN MAINE.

About the middle of May, 1917, a salmonid was caught in the tidal portion of the Penobscot River, Maine, which Mr. James D. De Rocher, superintendent of the Craig Brook fisheries station, sent to the Bureau of Fisheries for identification. At the time it was provisionally identified as *Salmo trutta* or *S. fario*. A recent examination of the specimen and comparison of it with *S. fario* indicates that it is quite positively the form that has been cultivated by the Bureau of Fisheries under the name of Scotch Sea Trout, commonly designated as *Salmo trutta*.

The total length of the specimen is 443 mm. and, as determined by an examination of the scales, it was probably between two and three years old when caught. It was a gravid female, near spawning condition—a curious fact for the time of the year, as the species usually spawns in the fall like other eastern Salmonidae.

The color in preservative is dusky on top of head and on back, sides silvery and belly white; many small, irregular black spots on sides of body, particularly above the lateral line, but some below posteriorly; spots of various shapes, some x-shaped and double-x-shaped, some irregularly ring-like. Their shape is largely determined by the margins of the

scales occupied. A few larger black spots are on the upper part of the opercle. The dorsal fin has rows of black spots, but all other fins are plain—pectoral with dusky tip; ventral and anal, pale; caudal, dusky and pale olive. Corroborative evidence that it is a so-called Scotch sea trout is the age as indicated by the scales, and that the only European salmonid planted in Maine coastal waters by the Bureau of Fisheries at a time to produce a fish of that age was the plant of the fiscal year, 1915, when 3,896 fry were placed in Alamoosook River, and 19,462 fry in Toddy Pond. Probably the present specimen originated in the Alamoosook River plant. Alamoosook River is a tributary to the tidal portion of Penobscot River, near Bucksport.

Br. 11-12; Gill-rakers  $7 + 12$  and  $7 + 12$ ; scales 29-120 + 4-29; on caudal peduncle downward and forward to lateral line 14, and from lateral line to anal, 14. D.  $3/9$ ; A.  $2/8$ ; P. 13; V. 9.

The following measurements are in percentages of the length from tip of snout to base of caudal fin, which is 391 mm.:

Distance from tip of snout to nape .....	14.8
Distance from nape to front of dorsal fin .....	34.8
Length of base of dorsal fin .....	12.8
Height of dorsal fin .....	13.6
Distance from posterior end of dorsal base to front base of adipose fin .....	25.8
Length of base of adipose .....	2.9
Distance from posterior end of adipose base to upper base of caudal .....	11.5
Distance from tip of snout to base of pectoral fin .....	21.2
Length of pectoral fin (frayed) .....	14.3?
Distance from base of pectoral fin to front of anal fin .....	35.8
Length of ventral fin (frayed) .....	11.5?
Distance from base of ventral fin to front of anal fin .....	23.8
Length of base of anal fin .....	8.9
Height of anal fin .....	13.3
Distance from posterior end of base of anal fin to lower base of caudal .....	14.1
Least depth of caudal peduncle .....	9.2
Greatest depth of body .....	24.3
Length of head .....	22.8

The following measurements are in percentages of the length of the head; which is 89 mm.:

Distance from tip of snout to posterior edge of preopercle	80.9
Vertical height of head in line through eye from lower edge of lower jaw to middle of interorbital space.....	51.6
Interorbital width .....	31.4
Length of eye .....	15.6
Length of snout .....	30.3
Distance from tip of snout to posterior end of maxillary.....	51.6
Length of maxillary .....	40.4
Length of lower jaw .....	61.7

WILLIAM CONVERSE KENDALL.

*U. S. Bureau of Fisheries.*

## REPTILES COLLECTED IN THE VICINITY OF CURRANT, NYE COUNTY, NEVADA.

Currant in Nye County, Nevada, is an isolated farming community bordering a small stream, Currant Creek, which has running water throughout the year. It lies on the western and somewhat southern slopes of the White Pine Ridge and ends some fifteen or twenty miles below in the desert. A heavy growth consisting principally of low willows borders the stream. There are large and tall Cottonwood trees in sheltered and swampy places. The valley is long and canyon-like, very narrow in places but widening enough in others to allow long, level strips for alfalfa, grains and small orchards. The hillsides when not rolling are high, steep, rocky and jagged, being sparsely covered with the characteristic desert flora such as sage-brush (*Artemisia tridentata*), rabbit-brush (*Chrysothamnus*), a few cacti (*Opuntia*), loco-weed (*Astragalus*), etc. It was on these rocky hillsides that most of the following reptiles were collected. The altitude is about 7,000 feet.

The reptiles were collected by myself in the spring of 1916. Some color notes were made of freshly killed specimens. The laboratory work was